

Table 10a. Numbers of Deaths¹ from or with Pneumoconiosis by State and U.S., 2000

	CA	CT	MA	ME	MI	NC	NE	NJ	NM ¹	NY	OR	WA	WI	US
All pneumoconiosis	143	24	42	<5	41	63	5	98	16	79	10	70	21	2,864
Coal workers' pneumoconiosis deaths	8	<5	0	0	9	9	0	<5	6	<5	<5	<5	<5	950
Asbestosis	122	20	40	<5	25	41	5	88	6	61	9	68	12	1,493
Silicosis	8	<5	<5	0	8	6	0	<5	<5	9	<5	<5	6	152
Other and unspecified pneumoconiosis	5	<5	<5	0	0	8	0	6	<5	5	<5	<5	<5	307

¹The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.

Table 10b. Age-Standardized Mortality Rate¹ from or with Selected Pneumoconiosis by State and U.S., 2000

	CA	CT	MA	MI	NC	NE	NJ	NM	NY	OR	WA	WI	US
Coal workers' pneumoconiosis	0.3	-- ²		1.2	1.6			4.8					4.4
Silicosis	0.3			1.1	1.0				0.6			1.4	0.7
Other and unspecified pneumoconiosis	0.2				1.4		0.9		0.3				1.4

¹Rates are expressed as deaths per one million residents

² NOTE: Dashes indicate that mortality rates were not calculated for states with less than five deaths.

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis by State and U.S., 2000

