

Table 10a. Numbers of Deaths from or with Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2001

	CA	CT	FL	LA	MA	ME	MI	NC	NE	NJ	NM	NY	OR	WA	WI	US
All pneumoconiosis	130	9	100	38	39	12	43	69	8	103	<5	88	9	69	27	2,747
Coal workers' pneumoconiosis deaths	<5	<5	10	<5	0	<5	7	8	<5	<5	0	11	<5	<5	<5	889
Asbestosis	113	6	80	31	33	12	32	49	7	96	<5	67	6	67	19	1,454
Silicosis	9	<5	<5	<5	<5	<5	<5	5	0	5	<5	<5	<5	<5	6	164
Other and unspecified pneumoconiosis	5	0	7	<5	<5	<5	<5	8	0	<5	0	6	<5	<5	<5	277

The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.


Table 10b. Age-Standardized Mortality Rates from or with Selected Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2001

	CA	CT	FL	LA	MA	ME	MI	NC	NE	NJ	NM	NY	OR	WA	WI	US
Coal workers' pneumoconiosis	--	--	0.6	--	--	--	0.9	1.4	--	--	--	0.7	--	--	--	4.0
Silicosis	0.3	--	--	--	--	--	--	0.8	--	0.7	--	--	--	--	1.4	0.7
Other and unspecified pneumoconiosis	0.2	--	0.4	--	--	--	--	1.4	--	--	--	0.4	--	--	--	1.2

Rates are expressed as deaths per one million residents

NOTE: Dashes indicate that mortality rates were not calculated for pneumoconiosis types with less than five deaths.

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis Ages 15 Years and Older by State and U.S., 2001



Rates are not illustrated for New Mexico because it had less than five total pneumoconiosis deaths.

Data Sources: Numbers of deaths: State Vital Records. Numbers of deaths in the U.S.: [National Center for Health Statistics](#) multiple cause of death file. Population statistics used to calculate rates: [U.S. Census Bureau](#)