

Table 10a. Numbers of Deaths¹ from or with Pneumoconiosis by State and U.S., 2002

	CA	CT	FL	LA	MA	ME	MI	NC	NJ	NM	NY	OR	WA	WI	US
All pneumoconiosis	131	26	105	45	36	17	47	46	89	10	74	12	79	26	2,718
Coal workers' pneumoconiosis deaths	7	0	13	5	0	<5	8	8	<5	5	5	0	>5	<5	858
Asbestosis	113	23	83	35	34	17	29	32	84	<5	58	11	73	19	1,473
Silicosis	<5	<5	6	<5	0	<5	<5	<5	<5	<5	9	0	<5	6	148
Other and unspecified pneumoconiosis	7	<5	<5	<5	<5	<5	6	5	<5	<5	<5	<5	0	<5	264

¹ The sum of these numbers may be greater than the total because cases could die with more than one type of pneumoconiosis listed for an individual death.

Table 10b. Age-Standardized Mortality Rate^{1,2} from or with Pneumoconiosis by State and U.S., 2002

	CA	CT	FL	LA	MA	ME	MI	NC	NJ	NM	NY	WA	WI	US
Coal workers' pneumoconiosis	0.3	N/A	0.7	1.5	N/A	N/A	1.0	1.3	N/A	4.2	0.3	0.8	0.2	3.8
Silicosis	N/A	N/A	0.3	N/A	N/A	N/A	N/A	N/A	N/A	N/A	1.0	0.4	1.4	0.7
Other and unspecified pneumoconiosis	0.3	N/A	N/A	N/A	N/A	N/A	0.7	0.8	N/A	N/A	N/A	N/A	0.2	1.1

¹Deaths per one million residents

² Rates were not calculated for states with less than five deaths in a category

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis by State and U.S., 2002

