

**Table 10a. Numbers of Deaths from or with Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2003**

	CA	CT	FL	KY	LA	MA	ME	MI	NC	NJ	NM	NY	OK	OR	WA	WI	US
All pneumoconiosis	138	22	101	67	45	36	12	45	62	81	13	73	12	14	71	32	2,639
Coal workers' pneumoconiosis	<5	0	17	49	<5	0	<5	<5	<5	<5	5	<5	<5	<5	<5	<5	773
Asbestosis	127	22	75	6	37	35	10	35	49	72	<5	61	9	13	70	22	1,471
Silicosis	6	<5	8	<5	<5	0	<5	<5	<5	5	<5	6	<5	<5	<5	8	179
Other and unspecified pneumoconiosis	<5	0	<5	10	<5	<5	<5	<5	6	<5	<5	<5	<5	<5	<5	0	241

The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.


**Table 10b. Age-Standardized Mortality Rates from or with Select Types of Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2003**

	CA	CT	FL	KY	LA	MA	ME	MI	NC	NJ	NM	NY	OK	OR	WA	WI	US
Coal workers' pneumoconiosis	—	—	0.9	15.4	—	—	—	—	—	—	3.8	—	—	—	—	—	3.4
Silicosis	0.3	—	0.3	—	—	—	—	—	—	0.7	—	0.4	—	—	—	1.8	0.8
Other and unspecified pneumoconiosis	—	—	—	3.1	—	—	—	—	1.0	—	—	—	—	—	—	—	1.1

Rates are expressed as deaths per one million residents.

NOTE: Dashes indicate that rates could not be calculated for pneumoconiosis types with less than five deaths.

**Figure 10. Age-Standardized Mortality Rates from or with Total Pneumoconiosis and Asbestosis, Ages 15 Years and Older, by State and U.S., 2003**



Data Sources: Numbers of deaths: State Vital Records. Numbers of deaths in the U.S.: [National Center for Health Statistics](http://www.cdc.gov/nchs/nvss/) multiple cause of death file. Population statistics used to calculate rates: [U.S. Census Bureau](http://www.census.gov/).