

Table 10a. Numbers of Deaths¹ from or with Pneumoconiosis by State and U.S., 2004

	CA	CT	FL	IA	KY	LA	MA	ME	MI	NJ	NM	NY	OK	OR	TX	WA	WI	US
All pneumoconiosis	123	21	108	11	72	31	42	16	45	76	8	84	13	11	147	72	33	2531
Coal workers' pneumoconiosis deaths	<5	<5	10	<5	50	<5	0	<5	<5	<5	<5	<5	<5	0	6	<5	0	703
Asbestosis	113	20	90	9	12	27	40	16	31	68	<5	65	8	9	128	71	20	1470
Silicosis	6	0	6	<5	<5	<5	0	<5	9	<5	<5	9	<5	<5	12	<5	12	166
Other and unspecified pneumoconiosis	<5	0	<5	<5	9	0	<5	<5	<5	<5	0	8	0	0	5	<5	<5	224

The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.


Table 10b. Age-Standardized Mortality Rate¹ from or with Pneumoconiosis by State and U.S., 2004

	CA	CT	FL	IA	KY	LA	MA	ME	MI	NJ	NM	NY	OK	OR	TX	WA	WI	US
Coal workers' pneumoconiosis	-	-	0.6	-	15.4	-	-	-	-	-	-	-	-	-	0.4	-	-	3.0
Silicosis	0.2	-	0.3	-	-	-	-	-	1.1	-	-	0.6	-	-	0.8	-	2.7	0.7
Other and unspecified pneumoconiosis	-	-	-	-	2.8	-	-	-	-	-	-	0.5	-	-	0.3	-	-	1.0

Rates are expressed as deaths per one million residents.

NOTE: Dashes indicate that rates could not be calculated for pneumoconiosis types with less than five deaths.

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis by State and U.S., 2004



Data Sources: Numbers of deaths: State Vital Records. Numbers of deaths in the U.S.: [National Center for Health Statistics](#) multiple cause of death file.

Population statistics used to calculate rates: [U.S. Census Bureau](#).